



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

JULY 3.

Mr. THOMAS MEEHAN, Vice-President, in the chair.

Ten persons present.

Note on Mazapilite, a new species.—Prof. GEO. A. KOENIG announced the occurrence of this mineral at Zacatecas, Mexico, in the mineral district of Mazapil. The crystals are well developed in all directions. They are of orthorhombic symmetry exhibiting a flat prism in combination with a brachy dome and a pyramid. The color is deep brown red, nearly black, but transparent at the edges. The hardness is nearly 7, its streak greenish yellow. The specific gravity 3.567. In closed tube a white crystalline sublimate is produced (As^2O^3) and water, while the powder turns dark brown. B. B. fuses at 3 to a black globule. On charcoal the odor of arsenic is observed. With borax only iron reaction. Easily soluble in in warm HCl. A preliminary analysis proves the mineral to be a *calcium ferric arsenite*. The structural formula must be made the subject of a more thorough investigation, which the speaker proposes to carry out in the fall. This mineral is the first representative of the class of pure arsenites in nature and is therefore of marked interest. For the material the author is indebted to the indefatigable zeal of Dr. F. A. Foote, who is now in Mexico.

The following was ordered to be printed:—